

## Tutorial Week 6

**Topics:** Cauchy sequences, completeness, uniform continuity.

**6.1.** Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be uniformly continuous functions between metric spaces. Prove that  $g \circ f: X \rightarrow Z$  is uniformly continuous.

*Solution.* Let  $\epsilon$  be a positive real number. The uniform continuity of  $f$  and  $g$  implies that there exists a positive real number  $\delta$  such that  $d_Y(y_1, y_2) < \delta$  implies  $d_Z(g(y_1), g(y_2)) < \epsilon$ , and there exists a positive real number  $\gamma$  such that  $d_X(x_1, x_2) < \gamma$  implies  $d_Y(f(x_1), f(x_2)) < \delta$ . Hence  $d_X(x_1, x_2) < \gamma$  implies  $d_Z((g \circ f)(x_1), (g \circ f)(x_2)) < \epsilon$ , and therefore  $g \circ f$  is uniformly continuous.  $\square$

**6.2.** Let  $S$  be a subset of a metric space  $(X, d_X)$  and let  $d_S$  be the induced metric on  $S$ .

- (a) Prove that the inclusion function  $\iota_S: S \rightarrow X$  is uniformly continuous.
- (b) Prove that a function  $f: (Y, d_Y) \rightarrow (S, d_S)$  is uniformly continuous if and only if  $\iota_S \circ f$  is uniformly continuous.

*Solution.*

- (a) Let  $\epsilon$  be a positive real number. Put  $\delta = \epsilon$ . If elements  $x_1$  and  $x_2$  of  $S$  satisfy  $d_S(x_1, x_2) < \delta$ , then

$$d_X(\iota_S(x_1), \iota_S(x_2)) = d_S(x_1, x_2) < \epsilon.$$

Hence  $\iota_S$  is uniformly continuous.

- (b) If  $f$  is uniformly continuous, then  $\iota_S \circ f$  is uniformly continuous because of part (a) and [Question 6.1](#).

Conversely, suppose  $\iota_S \circ f$  is uniformly continuous. Let  $\epsilon$  be a positive real number. Pick a positive real number  $\delta$  such that  $d_Y(y_1, y_2) < \delta$  implies  $d_X((\iota_S \circ f)(y_1), (\iota_S \circ f)(y_2)) < \epsilon$ . It follows that  $d_Y(y_1, y_2) < \delta$  implies

$$d_S(f(y_1), f(y_2)) = d_X((\iota_S \circ f)(y_1), (\iota_S \circ f)(y_2)) < \epsilon.$$

Hence  $f$  is uniformly continuous.  $\square$

**6.3.** Let  $(X, d_X)$ ,  $(Y, d_Y)$ , and  $(Z, d_Z)$  be metric spaces and let  $d$  be a metric on  $Y \times Z$  such that

$$\max\{d_Y(y_1, y_2), d_Z(z_1, z_2)\} \leq d((y_1, z_1), (y_2, z_2)) \leq d_Y(y_1, y_2) + d_Z(z_1, z_2)$$

for every pair of points  $(y_1, z_1)$  and  $(y_2, z_2)$  in  $Y \times Z$ .

- (a) Prove that the projections  $\pi_Y: Y \times Z \rightarrow Y$  and  $\pi_Z: Y \times Z \rightarrow Z$  are uniformly continuous.
- (b) Prove that a function  $f: X \rightarrow Y \times Z$  is uniformly continuous if and only if both  $\pi_Y \circ f$  and  $\pi_Z \circ f$  are.

*Solution.*

- (a) If  $\epsilon$  is a positive real number, then  $d((y_1, z_1), (y_2, z_2)) < \epsilon$  implies

$$d_Y(\pi_Y(y_1, z_1), \pi_Y(y_2, z_2)) = d_Y(y_1, y_2) \leq d((y_1, z_1), (y_2, z_2)) < \epsilon$$

and similarly  $d(\pi_Z(y_1, z_1), \pi_Z(y_2, z_2)) < \epsilon$ . Hence  $\pi_Y$  and  $\pi_Z$  are uniformly continuous.

- (b) If  $f$  is uniformly continuous, then it follows from [Question 6.1](#) and part (a) that both  $\pi_Y \circ f$  and  $\pi_Z \circ f$  are uniformly continuous.

Conversely, suppose both  $\pi_Y \circ f$  and  $\pi_Z \circ f$  are uniformly continuous. Let  $\epsilon$  be a positive real number. It follows from the uniform continuity of  $\pi_Y \circ f$  and  $\pi_Z \circ f$  that there exist positive real numbers  $\delta_Y$  resp.  $\delta_Z$  such that  $d_X(x_1, x_2) < \delta_Y$ , resp.  $d_X(x_1, x_2) < \delta_Z$  imply

$$d_Y((\pi_Y \circ f)(x_1), (\pi_Y \circ f)(x_2)) < \epsilon/2 \quad \text{resp.} \quad d_Z((\pi_Z \circ f)(x_1), (\pi_Z \circ f)(x_2)) < \epsilon/2.$$

Let  $\delta = \min\{\delta_Y, \delta_Z\}$ . It follows that  $d_X(x_1, x_2) < \delta$  implies

$$d(f(x_1), f(x_2)) \leq d_Y((\pi_Y \circ f)(x_1), (\pi_Y \circ f)(x_2)) + d_Z((\pi_Z \circ f)(x_1), (\pi_Z \circ f)(x_2)) < \epsilon,$$

so  $f$  is uniformly continuous. □

**6.4.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and let  $d$  be the sup norm metric on  $X \times Y$ .

- (a) Prove that the sequence  $((x_n, y_n))$  is Cauchy in  $X \times Y$  if and only if  $(x_n)$  is Cauchy in  $X$  and  $(y_n)$  is Cauchy in  $Y$ .
- (b) Prove that if  $X$  and  $Y$  are complete then  $X \times Y$  is complete. Is the converse true?

*Solution.*

- (a) Suppose  $((x_n, y_n))$  is a Cauchy sequence in  $(X \times Y, d)$ . By part (a) of [Question 6.3](#), both projections  $\pi_X: X \times Y \rightarrow X$  and  $\pi_Y: X \times Y \rightarrow Y$  are uniformly continuous. Hence  $(x_n) = (\pi_X(x_n, y_n))$  and  $(y_n) = (\pi_Y(x_n, y_n))$  are Cauchy because of [Proposition 2.60](#).

Conversely, suppose  $(x_n)$  is Cauchy in  $X$  and  $(y_n)$  is Cauchy in  $Y$ . Fix  $\epsilon > 0$ . Let  $N_x \in \mathbf{N}$  be such that for all  $m, n \geq N_x$  we have  $d_X(x_m, x_n) < \epsilon$ . Let  $N_y \in \mathbf{N}$  be such that for all  $m, n \geq N_y$  we have  $d_Y(y_m, y_n) < \epsilon$ . Let  $N = \max\{N_x, N_y\}$ , then for all  $m, n \geq N$  we have

$$d((x_m, y_m), (x_n, y_n)) = \max\{d_X(x_m, x_n), d_Y(y_m, y_n)\} < \epsilon,$$

so  $((x_n, y_n))$  is Cauchy in  $X \times Y$ .

- (b) Let  $((x_n, y_n))$  be a Cauchy sequence in  $X \times Y$ . By part (a),  $(x_n)$  is Cauchy in  $X$  and  $(y_n)$  is Cauchy in  $Y$ . Since  $X$  and  $Y$  are complete, we have  $(x_n) \rightarrow x \in X$  and  $(y_n) \rightarrow y \in Y$ . By [Question 5.4](#),  $((x_n, y_n)) \rightarrow (x, y) \in X \times Y$ .

The converse also holds: suppose  $X \times Y$  is complete. Let  $(x_n)$  be a Cauchy sequence in  $X$ , and fix some  $y \in Y$ . Then by (a) we have that  $((x_n, y))$  is Cauchy in  $X \times Y$ , so  $((x_n, y)) \rightarrow (x, y) \in X \times Y$ , which by [Question 5.4](#) implies that  $(x_n) \rightarrow x \in X$ . The same proof gives us that  $Y$  is complete. □

**6.5.** Suppose  $f: X \rightarrow Y$  is a *uniform homeomorphism* between metric spaces; that is, a homeomorphism such that both  $f$  and its inverse are uniformly continuous.

- (a) Prove that a sequence  $(x_n)$  is Cauchy in  $X$  if and only if  $(f(x_n))$  is Cauchy in  $Y$ .
- (b) Prove that  $X$  is complete if and only if  $Y$  is complete.
- (c) Prove that  $f: \mathbf{R} \rightarrow (-\pi/2, \pi/2)$  given by  $f(x) = \arctan(x)$  is uniformly continuous and a homeomorphism, but it is not a uniform homeomorphism.

- (d) Do you feel strongly that uniformly continuous functions ought to preserve completeness? (After all, they preserve Cauchy sequences, and completeness is defined in terms of Cauchy sequences.)

Prove that the function  $f$  defined in part (c) does not preserve completeness though it is uniformly continuous and a homeomorphism.

*Solution.* Let  $g: Y \rightarrow X$  denote the inverse of  $f$ .

- (a) By [Proposition 2.60](#),  $(x_n)$  being Cauchy implies  $(f(x_n))$  is Cauchy, while  $(f(x_n))$  being Cauchy implies  $(x_n) = (g(f(x_n)))$  is Cauchy.
- (b) Since  $X$  and  $Y$  are interchangeable, it suffices to prove one direction. Suppose  $X$  is complete and  $(y_n)$  is a Cauchy sequence in  $Y$ . It follows that  $(g(y_n))$  is Cauchy in  $X$ , and therefore converges to some point  $x$  in  $X$ . By [Theorem 2.52](#),  $(y_n) = (f(g(y_n)))$  converges to  $f(x)$ . Hence  $Y$  is complete.
- (c) Since  $f$  has an inverse  $\tan: (-\pi/2, \pi/2) \rightarrow \mathbf{R}$ , and both  $f$  and  $\tan$  are continuous. Hence  $f$  is a homeomorphism.

Given  $x_1 < x_2$ , apply the Mean Value Theorem to  $f(x) = \arctan(x)$  on  $[x_1, x_2]$  to get some  $\xi \in (x_1, x_2)$  such that

$$|f(x_2) - f(x_1)| = |f'(\xi)| |x_2 - x_1| = \frac{1}{1 + \xi^2} |x_2 - x_1| \leq |x_2 - x_1|.$$

So for any  $\varepsilon > 0$  we can take  $\delta = \varepsilon$  and conclude that  $f$  is uniformly continuous.

However, its inverse  $\tan: (-\pi/2, \pi/2) \rightarrow \mathbf{R}$  is not uniformly continuous, because  $(-\pi/2, \pi/2)$  is totally bounded (since bounded in  $\mathbf{R}$ ), but  $\mathbf{R}$  is not totally bounded. (Use [Proposition 2.71](#).)

- (d) The codomain  $(-\pi/2, \pi/2)$  is not complete because  $(\pi/2 - 1/n)$  is Cauchy but does not converge in  $(-\pi/2, \pi/2)$ . However, the domain  $\mathbf{R}$  is complete.  $\square$

**6.6.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f: X \rightarrow Y$  a surjective continuous function. Suppose that  $X$  is complete and for all  $x_1, x_2 \in X$  we have

$$d_X(x_1, x_2) \leq d_Y(f(x_1), f(x_2)).$$

Prove that  $Y$  is complete.

In particular, distance-preserving maps preserve completeness.

*Solution.* Let  $(y_n)$  be a Cauchy sequence in  $Y$ . For each  $n \in \mathbf{N}$ , let  $x_n \in f^{-1}(y_n)$ . I claim that  $(x_n)$  is a Cauchy sequence in  $X$ . Fix  $\varepsilon > 0$ . Let  $N \in \mathbf{N}$  be such that for all  $m, n \geq N$  we have  $d_Y(y_m, y_n) < \varepsilon$ . Then for all  $m, n \geq N$  we have

$$d_X(x_m, x_n) \leq d_Y(f(x_m), f(x_n)) = d_Y(y_m, y_n) < \varepsilon,$$

so  $(x_n)$  is indeed Cauchy in  $X$ .

Since  $X$  is complete, we have  $(x_n) \rightarrow x \in X$ , so that by the continuity of  $f$  we conclude that  $(y_n) = (f(x_n)) \rightarrow f(x) \in Y$ .  $\square$

**6.7.** Let  $(X, d)$  be a metric space.

- (a) Fix an arbitrary element  $y \in X$  and consider the function  $f: X \rightarrow \mathbf{R}$  given by  $f(x) = d(x, y)$ . Prove that  $f$  is uniformly continuous.

- (b) Prove that  $d: X \times X \rightarrow \mathbf{R}$  is uniformly continuous with respect to the sup metric  $D$  on  $X \times X$ .

*Solution.*

- (a) Let  $\varepsilon > 0$ . Set  $\delta = \varepsilon$ . If  $x, x' \in X$  satisfy  $d(x, x') < \delta = \varepsilon$ , then

$$|f(x) - f(x')| = |d(x, y) - d(x', y)| \leq d(x, x') < \varepsilon.$$

- (b) Let  $\varepsilon > 0$ . By part (a), there exists positive real numbers  $\delta_1$  and  $\delta_2$  such that  $d(x_1, x'_1) < \delta_1$  and  $d(x_2, x'_2) < \delta_2$  imply

$$d_{\mathbf{R}}(d(x_1, x_2), d(x'_1, x_2)) < \varepsilon/2 \quad \text{and} \quad d_{\mathbf{R}}(d(x'_1, x_2), d(x'_1, x'_2)) < \varepsilon/2.$$

Set  $\delta = \min\{\delta_1, \delta_2\}$ . If  $(x_1, x_2), (x'_1, x'_2) \in X \times X$  satisfy

$$\max\{d(x_1, x'_1), d(x_2, x'_2)\} = D((x_1, x_2), (x'_1, x'_2)) < \delta$$

then

$$d_{\mathbf{R}}(d(x_1, x_2), d(x'_1, x'_2)) \leq d_{\mathbf{R}}(d(x_1, x_2), d(x'_1, x_2)) + d_{\mathbf{R}}(d(x'_1, x_2), d(x'_1, x'_2)) < \varepsilon.$$

Hence  $d$  is uniformly continuous. □