

## Tutorial Week 9

**Topics:** sequence spaces, series.

**9.1** (Hölder's Inequality for  $\ell^1$  and  $\ell^\infty$ ). Prove that if  $u = (u_n) \in \ell^\infty$  and  $v = (v_n) \in \ell^1$ , then

$$\sum_{n=1}^{\infty} |u_n v_n| \leq \|u\|_{\ell^\infty} \|v\|_{\ell^1}.$$

*Solution.* Very straightforward.

By the definition of  $\ell^\infty$  and the  $\ell^\infty$ -norm, we have  $|u_n| \leq \|u\|_{\ell^\infty}$  for all  $n \in \mathbf{N}$ . Therefore for any  $m \in \mathbf{N}$  we have

$$\sum_{n=1}^m |u_n v_n| \leq \|u\|_{\ell^\infty} \sum_{n=1}^m |v_n|,$$

but the latter series converges because  $v \in \ell^1$ , to  $\|v\|_{\ell^1}$  and we get

$$\sum_{n=1}^{\infty} |u_n v_n| \leq \|u\|_{\ell^\infty} \|v\|_{\ell^1}. \quad \square$$

**9.2.** Prove that the norms on the sequence spaces  $\ell^\infty$  and  $\ell^p$  for  $p \neq 2$  cannot be defined by inner products.

*Solution.* We will verify that none of the norms in the question satisfy the Parallelogram Law (Proposition 3.12), so they cannot be defined by inner products. Consider  $e_1 = (1, 0, 0, 0, \dots)$  and  $e_2 = (0, 1, 0, 0, \dots)$ . Then

$$\|e_1 + e_2\|_{\ell^\infty}^2 + \|e_1 - e_2\|_{\ell^\infty}^2 = 2 \neq 4 = 2(\|e_1\|_{\ell^\infty}^2 + \|e_2\|_{\ell^\infty}^2)$$

and

$$\|e_1 + e_2\|_{\ell^p}^2 + \|e_1 - e_2\|_{\ell^p}^2 = 2 \times 2^{2/p} \neq 4 = 2(\|e_1\|_{\ell^p}^2 + \|e_2\|_{\ell^p}^2). \quad \square$$

**9.3.** Suppose  $1 \leq p \leq q$ . Prove that

$$\ell^p \subseteq \ell^q.$$

Show that if  $p < q$  then the inclusion is strict:  $\ell^p \subsetneq \ell^q$ .

*Solution.* We prove that

$$\|x\|_{\ell^q} \leq \|x\|_{\ell^p} \quad \text{for all } x \in \ell^p.$$

If  $\|x\|_{\ell^p} = 0$  then  $x = 0$  so  $\|x\|_{\ell^q} = 0$  and the inequality obviously holds. So suppose  $x \neq 0$ , then by dividing through by  $\|x\|_{\ell^p}$  we can reduce to proving that

$$\|x\|_{\ell^q} \leq 1 \quad \text{for all } x \text{ such that } \|x\|_{\ell^p} = 1.$$

But if  $\|x\|_{\ell^p} = 1$  then

$$\sum_{n=1}^{\infty} |x_n|^p = 1,$$

which means that for all  $n \in \mathbf{N}$  we have  $|x_n|^p \leq 1$ , so  $|x_n| \leq 1$ . However,  $p \leq q$  and  $|x_n| \leq 1$  implies that  $|x_n|^q \leq |x_n|^p$  for all  $n \in \mathbf{N}$ , so that

$$\|x\|_{\ell^q}^q = \sum_{n=1}^{\infty} |x_n|^q \leq \sum_{n=1}^{\infty} |x_n|^p = 1.$$

If  $p < q$  then  $\alpha := q/p > 1$ . For each  $n \in \mathbf{N}$ , let

$$x_n = \frac{1}{n^{1/p}},$$

so that

$$|x_n|^p = \frac{1}{n}, \quad |x_n|^q = \frac{1}{n^\alpha}.$$

We have

$$\|(x_n)\|_{\ell^p} = \sum_{n=1}^{\infty} \frac{1}{n} = \infty, \quad \|(x_n)\|_{\ell^q} = \sum_{n=1}^{\infty} \frac{1}{n^\alpha} < \infty,$$

so  $(x_n) \in \ell^q \setminus \ell^p$ . □

**9.4.** Prove that every finite-dimensional normed vector space is separable.

*Solution.* Let  $V$  be a finite-dimensional normed vector space. If  $V$  is a complex normed vector space, then we can restrict the scalars to real numbers to make  $V$  a real normed vector space. As a real vector space  $V$  is still finite-dimensional because  $\dim_{\mathbf{C}} V = 2 \dim_{\mathbf{R}} V$ . Hence, without loss of generality, we assume  $V$  is a real normed vector space.

Let  $v_1, \dots, v_n$  be a basis of  $V$ . By [Theorem 3.6](#), the norm on  $V$  is equivalent to the norm  $\|\cdot\|_1$  defined by

$$\|\alpha_1 v_1 + \dots + \alpha_n v_n\|_1 = |\alpha_1| + \dots + |\alpha_n|.$$

Since separability is a property of topological spaces and equivalent norms give rise to the same topology (see [Exercise 3.2](#)), we can assume without loss of generality that  $V$  is equipped with the norm  $\|\cdot\|_1$ .

Let  $W$  be the  $\mathbf{Q}$ -vector space spanned by  $\{v_1, \dots, v_n\}$ . We claim that  $V = \overline{W}$ .

If  $v$  is a vector in  $V$ , then there exists real numbers  $\alpha_1, \dots, \alpha_n$  such that  $v = \sum_{i=1}^n \alpha_i v_i$ . For each  $i$ , there exists a sequence of real numbers  $(a_{ij})$  converging to  $\alpha_i$  as  $j \rightarrow \infty$ . Define  $v_j = \sum_{i=1}^n a_{ij} v_i$ . We now prove  $v_j \rightarrow v$  as  $j \rightarrow \infty$ . If  $\epsilon$  is a positive real number, then for every index  $i$  there exists a positive integer  $N_i$  such that  $n > N_i$  implies  $|\alpha_i - a_{ij}| < \epsilon/n$ . Put  $N = \max\{N_1, \dots, N_n\}$ . It follows that  $n > N$  implies

$$\|v - v_j\| = \sum_i |\alpha_i - a_{ij}| < n \cdot (\epsilon/n) = \epsilon.$$

The space  $W$  is countable because it is the Cartesian product of  $n$  copies of  $\mathbf{Q}$ . Hence  $W$  is a countable dense subset of  $V$ . □

**9.5.** Let  $c_{00}$  be the space of sequences with only finitely many nonzero terms (see [Example 3.23](#)), and consider it as a subspace of  $\ell^\infty$ . Prove that  $c_{00}$  is separable.

*Solution.* Let  $e_j$  be the sequence whose  $j$ -th entry is 1, and all the others are 0. Since every vector in  $c_{00}$  has only finitely many non-zero terms, it follows that  $c_{00}$  is spanned by  $\{e_1, e_2, \dots\}$ . Let  $V_n$  be the span of  $\{e_1, \dots, e_n\}$ , which is separable by [Question 9.4](#). Pick a countable dense subset  $S_n$  of  $V_n$  and put  $S = \bigcup_{n=1}^{\infty} S_n$ . We claim that  $S$  is a countable dense subset of  $c_{00}$ .

As a countable union of countable sets,  $S$  is countable because of [Exercise 1.2](#). If  $v$  is a vector in  $c_{00}$ , then  $v$  belongs to  $V_n$  for some  $n$ , so there exists a sequence in  $S_n$  converging to  $v$ , which is also a sequence in  $S$  converging to  $v$ . Hence  $S$  is dense in  $c_{00}$ .

*Alternative:* Once you see that  $c_{00}$  is spanned by the countable set  $\{e_1, e_2, \dots\}$ , use [Proposition 3.31](#). □

**9.6.** Consider the subset  $c_0 \subseteq \mathbf{F}^{\mathbf{N}}$  of all sequences with limit 0:

$$c_0 = \{(a_n) \in \mathbf{F}^{\mathbf{N}} : (a_n) \rightarrow 0\}.$$

- (a) Prove that  $c_0$  is a closed subspace of  $\ell^\infty$ .
- (b) Conclude that  $c_0$  is a Banach space.
- (c) Prove that  $c_0$  is separable.

*Solution.*

- (a) It's pretty clear that  $c_0$  is a subspace of  $\mathbf{F}^{\mathbf{N}}$ , and hence of  $\ell^\infty$ . To show that  $c_0$  is closed in  $\ell^\infty$ , let  $(x_n) \rightarrow x \in \ell^\infty$  with  $x_n \in c_0$  for all  $n \in \mathbf{N}$ . We want to prove that  $x \in c_0$ .

Write  $x_n = (a_{nm}) = (a_{n1}, a_{n2}, a_{n3}, \dots)$  and  $x = (a_m) = (a_1, a_2, a_3, \dots)$ . Let  $\varepsilon > 0$ . There exists  $N \in \mathbf{N}$  such that for all  $n \geq N$  we have

$$\sup_m |a_m - a_{nm}| = \|x - x_n\|_{\ell^\infty} < \frac{\varepsilon}{2}.$$

Consider the sequence  $x_N = (a_{Nm}) \in c_0$ . It converges to 0, so that there exists  $M \in \mathbf{N}$  such that for any  $m \geq M$  we have

$$|a_{Nm}| < \frac{\varepsilon}{2}.$$

Therefore, for  $m \geq M$ , we get

$$|a_m| = |a_m - a_{Nm} + a_{Nm}| \leq |a_m - a_{Nm}| + |a_{Nm}| < \varepsilon.$$

Hence  $x = (a_m) \rightarrow 0$ .

- (b) Since  $c_0$  is closed and  $\ell^\infty$  is Banach,  $c_0$  is Banach.
- (c) I claim that  $c_0$  has the same Schauder basis as the one given in [Example 3.32](#) for  $\ell^p$ :  $\{e_1, e_2, \dots\}$  where  $e_n = (0, \dots, 0, 1, 0, \dots)$  with the 1 in the  $n$ -th spot. Take  $v = (v_n) \in c_0$ , then  $(v_n) \rightarrow 0$ . I claim that the series

$$\sum_{n=1}^{\infty} v_n e_n$$

converges to  $v$  with respect to the norm on  $c_0$ , which is the  $\ell^\infty$ -norm:

$$\left\| v - \sum_{n=1}^m v_n e_n \right\|_{\ell^\infty} = \|(0, \dots, 0, v_{m+1}, v_{m+2}, v_{m+3}, \dots)\|_{\ell^\infty} = \sup_{n \geq m+1} |v_n|,$$

and the latter converges to 0 as  $m \rightarrow \infty$ , since  $(v_n) \rightarrow 0$ . The uniqueness of the coefficients follows in precisely the same way as for [Example 3.32](#).  $\square$

**9.7.** Consider the space  $\ell^\infty$  of bounded sequences.

- (a) Let  $S \subseteq \ell^\infty$  be the subset of sequences  $(a_n)$  such that  $a_n \in \{0, 1\}$  for all  $n \in \mathbf{N}$ . Prove that  $S$  is an uncountable set.  
*[Hint: Mimic Cantor's diagonal argument.]*
- (b) Use  $S$  to construct an uncountable set  $T$  of disjoint open balls in  $\ell^\infty$ .
- (c) Conclude that  $\ell^\infty$  is not separable.

*Solution.*

(a) Suppose  $S$  is countable and enumerate its elements:

$$\begin{aligned} a_1 &= (a_{11}, a_{12}, a_{13}, \dots) \\ a_2 &= (a_{21}, a_{22}, a_{23}, \dots) \\ a_3 &= (a_{31}, a_{32}, a_{33}, \dots) \\ &\vdots \end{aligned}$$

Go down the diagonal of this infinite grid of 0's and 1's, and define  $b_n = 1 - a_{nn}$  for all  $n \in \mathbf{N}$ . Then  $b = (b_n) \in S$ , but  $b \neq a_m$  for any  $m \in \mathbf{N}$ , contradiction.

(b) If  $a = (a_n), b = (b_n) \in S$  with  $a \neq b$  then

$$\|a - b\| = \sup_n |a_n - b_n| = 1,$$

so  $\mathbf{B}_{1/2}(a) \cap \mathbf{B}_{1/2}(b) = \emptyset$ .

Therefore we can take

$$T = \{\mathbf{B}_{1/2}(s) : s \in S\}.$$

(c) Any dense subset  $D$  of  $\ell^\infty$  must contain at least one point (in fact, must be dense) in each open ball in the set  $T$ . Since  $T$  is uncountable,  $D$  must also be uncountable, so  $\ell^\infty$  is not separable.  $\square$

**9.8.** Give an example of a series that converges but does not converge absolutely.

*Solution.* In  $\mathbf{R}$ , consider

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n}.$$

Taking absolute values we get the harmonic series, which does not converge.

The original series has alternating signs, and its terms in absolute value form a decreasing sequence  $(1/n)$  that converges to zero, hence the series converges by the alternating series test.  $\square$

**9.9.** If a series  $\sum_{n=1}^{\infty} a_n$  in a normed space  $(V, \|\cdot\|)$  converges absolutely, then

$$\left\| \sum_{n=1}^{\infty} a_n \right\| \leq \sum_{n=1}^{\infty} \|a_n\|.$$

*Solution.* This follows from the usual triangle inequality.

For any  $m \in \mathbf{N}$ , we have

$$\|a_1 + \dots + a_m\| \leq \|a_1\| + \dots + \|a_m\|.$$

Taking limits as  $m \rightarrow \infty$  we get

$$\left\| \sum_{n=1}^{\infty} a_n \right\| = \left\| \lim_{m \rightarrow \infty} \sum_{n=1}^m a_n \right\| = \lim_{m \rightarrow \infty} \left\| \sum_{n=1}^m a_n \right\| \leq \lim_{m \rightarrow \infty} \sum_{n=1}^m \|a_n\| = \sum_{n=1}^{\infty} \|a_n\|. \quad \square$$