

Tutorial Week 10

Topics: continuous linear transformations, series, projections.

10.1. Prove that all linear transformations between finite-dimensional normed vector spaces are continuous.

Solution. Let v_1, \dots, v_m be a basis of V and let w_1, \dots, w_n be a basis of W . By [Theorem 3.6](#), the norms on V and on W are equivalent to the norms $\|\cdot\|_V$ and $\|\cdot\|_W$ defined by

$$\|\alpha_1 v_1 + \dots + \alpha_m v_m\|_V = |\alpha_1| + \dots + |\alpha_m| \quad \text{and} \quad \|\beta_1 w_1 + \dots + \beta_n w_n\|_W = |\beta_1| + \dots + |\beta_n|.$$

Since continuity of f is determined by the topologies on V and W , and since equivalent norms give rise to the same topology (see [Exercise 3.2](#)), we can assume without loss of generality that V and W are equipped with the norms $\|\cdot\|_V$ and $\|\cdot\|_W$ respectively.

Put $M = \max\{\|f(v_1)\|_W, \dots, \|f(v_m)\|_W\}$. If $v = \alpha_1 v_1 + \dots + \alpha_m v_m$, then

$$\begin{aligned} \|f(v)\|_W &= \|f(\alpha_1 v_1 + \dots + \alpha_m v_m)\|_W \\ &= \|\alpha_1 f(v_1) + \dots + \alpha_m f(v_m)\|_W \\ &\leq \|\alpha_1 f(v_1)\|_W + \dots + \|\alpha_m f(v_m)\|_W \\ &= |\alpha_1| \|f(v_1)\|_W + \dots + |\alpha_m| \|f(v_m)\|_W \\ &\leq \alpha_1 M + \dots + \alpha_m M \leq M \|v\|_V. \end{aligned}$$

Hence f is Lipschitz, and therefore continuous by [Proposition 3.25](#). □

10.2. Let $f_1: V \rightarrow W_1$ and $f_2: V \rightarrow W_2$ be two continuous linear transformations between normed vector spaces. Prove that the function $f: V \rightarrow W_1 \times W_2$ defined by $f(v) = (f_1(v), f_2(v))$ is a continuous linear transformation.

Solution. We start with proving that f is linear. If v_1 and v_2 are vectors in V , then

$$\begin{aligned} f(v_1 + v_2) &= (f_1(v_1 + v_2), f_2(v_1 + v_2)) \\ &= (f_1(v_1) + f_1(v_2), f_2(v_1) + f_2(v_2)) \\ &= (f_1(v_1), f_2(v_1)) + (f_1(v_2), f_2(v_2)) = f(v_1) + f(v_2). \end{aligned}$$

If α is a scalar and v is a vector in V , then

$$f(\alpha v) = (f_1(\alpha v), f_2(\alpha v)) = (\alpha f_1(v), \alpha f_2(v)) = \alpha f(v).$$

Hence f is linear.

It remains to prove that f is continuous. Let $\pi_1: W_1 \times W_2 \rightarrow W_1$ and $\pi_2: W_1 \times W_2 \rightarrow W_2$ be the projections. Since $\pi_1 \circ f = f_1$ and $\pi_2 \circ f = f_2$, it follows from [Question 3.7](#) that f is continuous. □

10.3. Let c_{00} be the space of sequences with only finitely many nonzero terms (see [Example 3.23](#)), which is considered as a subspace of ℓ^∞ . Let $f: c_{00} \rightarrow \mathbf{F}^{\mathbf{N}}$ be the function defined by $(f(v))_n = nv_n$.

- (a) Prove that the image of the function f is contained in ℓ^∞ .
- (b) Let $g: c_{00} \rightarrow \ell^\infty$ be the function defined by $g(v) = f(v)$. Prove that g is not continuous.
- (c) Prove that there exists a discontinuous linear transformation from ℓ^∞ to itself.

In this part, you can use the following fact:

Let V and W be \mathbf{F} -vector spaces. If S is a subspace of V and if $\phi: S \rightarrow W$ is a linear transformation, then there exists a linear transformation $\tilde{\phi}: V \rightarrow W$ such that $\phi = \tilde{\phi}|_S$.

Solution.

- (a) Let $v \in c_{00}$. By the definition of c_{00} , there exists a positive integer N such that $n > N$ implies $v_n = 0$. It follows that $(f(v))_n \leq Nv_n$ for every positive integer n . Hence $\|f(v)\|_{\ell^\infty} \leq N\|v\|_{\ell^\infty} < \infty$, which implies $f(v) \in \ell^\infty$.
- (b) For every positive integer n , let $v^{(m)}$ be the vector in ℓ^∞ defined by

$$v_n^{(m)} = \begin{cases} 1 & \text{if } n \leq m, \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to verify $\|v^{(m)}\|_{\ell^\infty} = 1$ but $\|g(v^{(m)})\|_{\ell^\infty} = m$ for every positive integer m . Hence g is not Lipschitz, which implies g is not continuous by [Proposition 3.25](#).

- (c) Apply the fact to the discontinuous linear transformation $g: c_{00} \rightarrow \ell^\infty$ in part (b) to obtain a linear transformation $\tilde{g}: \ell^\infty \rightarrow \ell^\infty$, which cannot be continuous because otherwise its restriction g would be continuous. \square

10.4. If $f \in L(V, W)$ with V, W normed spaces, and the series

$$\sum_{n=1}^{\infty} \alpha_n v_n, \quad \alpha_n \in \mathbf{F}, v_n \in V,$$

converges in V , then the series

$$\sum_{n=1}^{\infty} \alpha_n f(v_n)$$

converges in W to the limit

$$f\left(\sum_{n=1}^{\infty} \alpha_n v_n\right).$$

Solution. Let

$$x_m = \sum_{n=1}^m \alpha_n v_n, \quad x = \sum_{n=1}^{\infty} \alpha_n v_n.$$

We know that $(x_m) \rightarrow x$ in V .

Since $f \in L(V, W)$ is continuous, we have that $(f(x_m)) \rightarrow f(x)$ in W .

But f is also linear, so

$$f(x_m) = \sum_{n=1}^m \alpha_n f(v_n).$$

Hence

$$\left(\sum_{n=1}^m \alpha_n f(v_n)\right) \rightarrow f(x),$$

so that the series

$$\sum_{n=1}^{\infty} \alpha_n f(v_n) \quad \text{converges to } f(x). \quad \square$$

10.5.

(a) Prove that the function $f: \ell^1 \rightarrow \mathbf{F}$ defined by

$$f((a_n)) = \sum_{n=1}^{\infty} a_n.$$

is continuous.

(b) Prove that the following subset is a closed subspace of ℓ^1 :

$$S = \left\{ (a_n) \in \ell^1 : \sum_{n=1}^{\infty} a_n = 0 \right\}.$$

Solution.

(a) First note that this is a reasonable definition, because the infinite series on the right hand side converges in \mathbf{F} :

$$\left| \sum_{n=1}^N a_n \right| \leq \sum_{n=1}^N |a_n|,$$

and the latter converges as $N \rightarrow \infty$ since $(a_n) \in \ell^1$.

The function f is linear. It is also Lipschitz, because as we have just seen:

$$|f((a_n))| = \left| \sum_{n=1}^{\infty} a_n \right| \leq \sum_{n=1}^{\infty} |a_n| = \|(a_n)\|_{\ell^1}.$$

Hence f is continuous by [Proposition 3.25](#).

(b) As the kernel of the continuous linear functional f , the subset S of ℓ^1 is a closed subspace. □

10.6. Fix $j \in \mathbf{N}$ and consider the map $\pi_j: \mathbf{F}^{\mathbf{N}} \rightarrow \mathbf{F}$ given by

$$\pi_j((a_n)) = a_j.$$

(a) Show that π_j is linear.

(b) Prove that the restriction of π_j to ℓ^p for $1 \leq p \leq \infty$ is continuous and surjective.

Solution.

(a) Straightforward.

(b) We have for $a \in \ell^\infty$:

$$|\pi_j(a)| = |a_j| \leq \sup_{n \geq 1} \{|a_n|\} = \|a\|_{\ell^\infty},$$

so π_j is bounded.

Similarly for $a \in \ell^p$:

$$|\pi_j(a)| = |a_j| = (|a_j|^p)^{1/p} \leq \left(\sum_{n \geq 1} |a_n|^p \right)^{1/p} = \|a\|_{\ell^p}.$$

For the surjectivity we note that for any $a \in \mathbf{F}$ we have $\pi_j((0, \dots, 0, a, 0, \dots)) = a$ and $(0, \dots, 0, a, 0, \dots) \in \ell^1 \subseteq \ell^p$ for all $1 \leq p \leq \infty$. □

10.7. Let V be a normed space and φ, ψ be commuting projections: $\varphi \circ \psi = \psi \circ \varphi$. Prove that $\varphi \circ \psi$ is a projection with image $\text{im } \varphi \cap \text{im } \psi$.

Solution. We know that the composition of continuous linear maps is continuous linear, so this is true for $\varphi \circ \psi$. To conclude that it is a projection, we need to compute its square:

$$(\varphi \circ \psi) \circ (\varphi \circ \psi) = (\varphi \circ \varphi) \circ (\psi \circ \psi) = \varphi \circ \psi,$$

where it was crucial that φ and ψ commute.

For the statement about the image, note that $w \in \text{im}(\varphi \circ \psi)$ if and only if there exists $v \in V$ such that

$$w = \varphi(\psi(v)) = \psi(\varphi(v)),$$

which implies that $w \in \text{im} \varphi \cap \text{im} \psi$. So $\text{im}(\varphi \circ \psi) \subseteq \text{im} \varphi \cap \text{im} \psi$.

Conversely, suppose $w \in \text{im} \varphi \cap \text{im} \psi$, then there exists $v \in V$ such that $w = \psi(v)$. But $w \in \text{im} \varphi$ and φ is a projection, so that

$$w = \varphi(w) = \varphi(\psi(v)) \in \text{im}(\varphi \circ \psi). \quad \square$$

10.8. Let φ be a nonzero orthogonal projection (that is, φ is not the constant function 0) on an inner product space V . Prove that $\|\varphi\| = 1$.

Solution. We know that $(\text{im} \varphi)^\perp = \ker \varphi$. For any x we have

$$\varphi(x - \varphi(x)) = \varphi(x) - \varphi^2(x) = \varphi(x) - \varphi(x) = 0,$$

so $x - \varphi(x) \in \ker \varphi$. Therefore

$$\langle x, \varphi(x) \rangle - \|\varphi(x)\|^2 = \langle x - \varphi(x), \varphi(x) \rangle = 0,$$

so

$$\|\varphi(x)\|^2 = \langle x, \varphi(x) \rangle \leq \|x\| \|\varphi(x)\|$$

by the Cauchy–Schwarz Inequality. Hence $\|\varphi(x)\| \leq \|x\|$ for all x , hence $\|\varphi\| \leq 1$. However for $x \in \text{im} \varphi$ we have $\varphi(x) = x$ so $\|\varphi(x)\| = \|x\|$ and we conclude that $\|\varphi\| = 1$. \square