

# 2 Set theory

## Chapter contents

2.1	Sets . . . . .	2
2.2	Relations, functions, orders . . . . .	17
2.3	The real numbers . . . . .	26
2.4	Bounds . . . . .	29
2.5	Distance on the real line . . . . .	39

## 2.1 Sets

**Definition 2.1.** A *set* is a collection of unique objects, called *elements*. If  $A$  is a set and  $x$  is an element of  $A$ , we write  $x \in A$ . If  $x$  is not an element of  $A$ , we write  $x \notin A$ .

**Example 2.2.**  $A = \{\mathbf{Z}, 0, \mathbf{Q}, \mathbf{R}, \{\pi, e, 1\}\}$  is a set.

We have:  $\mathbf{Z} \in A$ ,  $\mathbf{N} \notin A$ ,  $0 \in A$ ,  $1 \notin A$ .

The elements of a set are unordered:

$$\{\mathbf{Z}, 0, \mathbf{Q}, \mathbf{R}, \{\pi, e, 1\}\} = \{0, \mathbf{Q}, \mathbf{Z}, \{\pi, e, 1\}, \mathbf{R}\}.$$

Most interesting sets are too big or too complicated to be described by a complete explicit enumeration of their elements.

**Example 2.3.** The set of multiples of 17:

$$S = \{x \in \mathbf{Z} : (\exists k \in \mathbf{Z})x = 17k\}$$

The set of Fermat primes:

$$S = \{x \in \mathbf{Z} : (\exists k \in \mathbf{Z})x = 2^{2^k} + 1 \text{ and } x \text{ is prime}\}$$

This type of description is called *set comprehension* and is close to the informal natural language description. It is very powerful, but with great power

**Example 2.4.** Let  $S$  be the set of all things that are not elephants.

Some elements of  $S$ :

If you don't see a problem with the above, consider

**Example 2.5** (Russell's Paradox). Let  $P$  be the set of all sets that are not elements of themselves. Is  $P \in P$ ?

**Example 2.6** (Barber Paradox). There is a barber who shaves all those, and only those, who do not shave themselves. Does the barber shave himself?

Issues such as these paradoxes forced mathematicians to develop formal axiomatic descriptions of set theory.

## ZFC axioms of set theory

The most commonly used formalisms are ZFC (Zermelo–Fraenkel with Choice) or one of its variants such as NBG (von Neumann–Bernays–Gödel).

**Definition 2.7** (ZFC axioms).

- **Extensionality:** two sets are equal if and only if they have the same elements.
- **Pairing:** for any  $a$  and  $b$  there exists a set  $\{a, b\}$  that contains exactly  $a$  and  $b$ .
- **Separation:** if  $U$  is a set and  $p(x)$  is a condition on  $U$  expressed in first-order logic, there exists a set  $X = \{x \in U : p(x)\}$ .
- **Union:** if  $A$  is a set of sets  $X_\alpha$ , there exists a set  $\bigcup_{\alpha \in A} X_\alpha$  that contains all elements that belong to at least one  $X_\alpha$ .
- **Power set:** if  $X$  is a set, there exists a set  $P(X)$  that contains all the subsets of  $X$ .
- **Inductive set:** there exists a set  $N$  such that every  $x \in N$  is a set,  $\emptyset \in N$ , and if  $x \in N$  then  $x \cup \{x\} \in N$ .
- **Replacement:** if  $f$  is a function, the range of  $f$  is a set.
- **Regularity:** if  $A$  is a non-empty set, there exists  $y \in A$  such that  $A$  and  $y$  have no common elements.
- **Choice:** if  $A$  is a set of nonempty sets  $X_\alpha$ , there exists a set that contains exactly one element from each set  $X_\alpha$ .

## Equality

The Axiom of Extensionality defines the notion of equality between two sets:

**Definition 2.8.** Two sets  $A$  and  $B$  are *equal* (we write:  $A = B$ ) when

$$(\forall x)x \in A \Leftrightarrow x \in B.$$

This suggests the possibility of a weaker notion:

**Definition 2.9.** A set  $A$  is a *subset* of a set  $B$  (we write:  $A \subseteq B$ ) when

$$(\forall x)x \in A \Rightarrow x \in B.$$

If  $A \subseteq B$  and  $A \neq B$  we write  $A \subsetneq B$  and say that  $A$  is a *proper subset* of  $B$ .

**Example 2.10.**  $X = \{1, 2, 3, 4, 5\}$ .

## Separation

The Axiom of Separation formalises the correct use of set comprehension: if  $U$  is a set and  $p(x)$  is a condition on  $U$ , there exists a set

$$X = \{x \in U : p(x) \text{ is True}\}$$

consisting precisely of the elements  $x$  of  $U$  for which  $p(x)$  is **True**.

**Example 2.11.** Take  $U = \{1, 2, 3, 4, 5\}$  and  $p(x) : x < 4$ .

Note that the Axiom of Separation implies the existence of the *empty set*  $\emptyset = \{\}$ , with no elements.

## Operations with sets

Here we assume that all the sets we consider are subsets of a fixed set  $U$ .

**Definition 2.12.** Let  $A$  and  $B$  be sets.

- the *union* of  $A$  and  $B$  is

$$A \cup B = \{x \in U : x \in A \text{ or } x \in B\}.$$


- the *intersection* of  $A$  and  $B$  is

$$A \cap B = \{x \in U : x \in A \text{ and } x \in B\}.$$

- the *difference* (sometimes called the *complement* of  $B$  in  $A$ ) is


$$A \setminus B = \{x \in U : x \in A \text{ and } x \notin B\}.$$

**Example 2.13** (Everybody ♥ Venn diagrams).



**Example 2.14** (Everybody ♥ Venn diagrams).

Let  $A = \{1, 2, 3\}$ ,  $B = \{2, 3, 4, 5\}$ ,  $C = \{3, 5, 6\}$ .



$$A \cup B =$$

$$A \cap B =$$

$$A \setminus B =$$

$$(A \cap C) \setminus B =$$

## Cartesian product

Let  $A$  and  $B$  be sets.

**Definition 2.15.** An *ordered pair*  $(a, b)$  is a set of the form

$$(a, b) = \{\{a\}, \{a, b\}\}, \quad \text{where } a \in A \text{ and } b \in B.$$

**Example 2.16.**  $(1, 2) = \{\{1\}, \{1, 2\}\}.$

The essential property of ordered pairs is

given  $a_1, a_2 \in A$  and  $b_1, b_2 \in B$ , we have  $(a_1, b_1) = (a_2, b_2)$  if and only if  $a_1 = a_2$  and  $b_1 = b_2$ .

Let  $A$  and  $B$  be sets.

**Definition 2.17.** The *Cartesian product* of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}.$$

**Example 2.18.** Let  $A = \{1, 5, \text{dog}\}$  and  $B = \{5, \text{cat}\}$ .

## Some proofs with sets

**Theorem 2.19.** *Let  $A$  and  $B$  be sets. We have  $A = B$  if and only if  $(A \subseteq B \text{ and } B \subseteq A)$ .*

**Theorem 2.20.** *Let  $A$ ,  $B$ , and  $C$  be sets. Then  $(B \setminus A) \cap C = (B \cap C) \setminus A$ .*

## Numbers

We spend lots of time working with sets of numbers. Recall the most common ones:

The *natural numbers*

$$\mathbf{N} = \{0, 1, 2, 3, 4, \dots\}.$$

Informally, we can think of this as:  $0 \in \mathbf{N}$ , and if  $n \in \mathbf{N}$  then  $n + 1 \in \mathbf{N}$ . This should remind you of the Principle of Mathematical Induction!

**Theorem 2.21** (Well-Ordering Property of  $\mathbf{N}$ ). *Every non-empty subset  $S \subseteq \mathbf{N}$  has a *minimum*: there exists  $m \in S$  such that  $m \leq x$  for all  $x \in S$ .*

Starting from the natural numbers  $\mathbf{N}$ , we successively broaden our numerical horizons:

- the *integers*

$$\mathbf{Z} = \{ \pm n : n \in \mathbf{N} \} = \{ \dots, -3, -2, -1, 0, 1, 2, 3, \dots \};$$

- the *rational numbers*

$$\mathbf{Q} = \left\{ \frac{a}{b} : a \in \mathbf{Z}, b \in \mathbf{Z}_{>0} \right\}, \quad \text{with } \frac{a}{b} = \frac{c}{d} \Leftrightarrow ad = bc;$$

- the *real numbers*  $\mathbf{R}$ , to be discussed in more depth soon;

- the *complex numbers*

$$\mathbf{C} = \{ a + ib : a \in \mathbf{R}, b \in \mathbf{R} \}, \quad \text{where } i^2 = -1.$$

We have  $\mathbf{N} \subsetneq \mathbf{Z} \subsetneq \mathbf{Q} \subsetneq \mathbf{R} \subsetneq \mathbf{C}$ .

We also have the standard addition and multiplication operations on these sets, with the usual arithmetic properties.

## 2.2 Relations, functions, orders

**Definition 2.22.** Let  $A$  and  $B$  be sets. A *relation* from  $A$  to  $B$  is a subset  $R \subseteq A \times B$ .

For  $a \in A$ ,  $b \in B$ , we write  $aRb$  if  $(a, b) \in R$ .

**Example 2.23.** The University of Melbourne has a database of the students enrolled in various subjects.

Let  $P$  be the set of all students and  $S$  the set of all subjects. Then the University's enrollment database can be described mathematically as

We will soon talk about two special types of relations that are important to analysis: functions and orders.

It is worth noting that other mathematical concepts can be naturally viewed as relations.

**Example 2.24** (Equality in a set). Let  $X$  be a set and consider the *diagonal subset* of  $X \times X$ , given by

$$\Delta(X) = \{(x, x) : x \in X\}$$

**Example 2.25** (Element of a set). Let  $X$  be a set and let  $P(X)$  be its power set. Consider

$$R = \{(x, A) \in X \times P(X) : x \in A\}.$$

**Example 2.26** (Subset). Let  $X$  be a set and let  $P(X)$  be its power set. Consider

$$R = \{(A, B) \in P(X) \times P(X) : A \subseteq B\}.$$

## Functions

**Definition 2.27.** Let  $A$  and  $B$  be sets; let  $f \subseteq A \times B$  be a relation from  $A$  to  $B$ .

We say that  $f$  is a *function* from  $A$  to  $B$  if: for every  $x \in A$  there exists a unique  $y \in B$  such that  $(x, y) \in f$ .

We write  $f : A \longrightarrow B$ ;  $A$  is called the *domain* of  $f$ ;  $B$  is called the *codomain* of  $f$ .

If  $(x, y) \in f$  then we write  $y = f(x)$ ;  $y$  is called the *image* of  $x$ . The set

$$f^{-1}(y) = \{x \in A : f(x) = y\} \subseteq A$$

is called the *preimage* of  $y$ .

More generally, if  $S \subseteq A$  then the set

$$f(S) = \{y \in B : (\exists x \in S) f(x) = y\} \subseteq B$$

is called the *image* of  $S$ . In the special case  $S = A$ , the set  $f(A)$  is called the *image* (or *range*) of  $f$ .

If  $T \subseteq B$  then the set

$$f^{-1}(T) = \{x \in A : f(x) \in T\} \subseteq A$$

is called the *preimage* of  $T$ .

**Example 2.28.** Let  $f : \mathbf{R} \longrightarrow \mathbf{R}$  be the function defined by  $f(z) = z^2$  for all  $z \in \mathbf{R}$ .

**Definition 2.29.** Let  $A$  and  $B$  be sets and let  $f : A \longrightarrow B$  be a function. We say that  $f$  is ...

- ... *injective* if  $(\forall x, y \in A) (f(x) = f(y) \Rightarrow x = y)$ ;
- ... *surjective* if  $(\forall z \in B) (\exists x \in A) f(x) = z$ ;
- ... *bijjective* if  $f$  is both injective and surjective.

**Example 2.30.** The function  $f : \mathbf{R} \longrightarrow \mathbf{R}$  given by  $f(x) = e^x$  is

**Example 2.31.** The function  $f : \mathbf{R} \longrightarrow \mathbf{R}_{\geq 0}$  given by  $f(x) = x^2$  is

**Example 2.32** (Identity function). For any set  $A$ , there exists a function  $\text{id}_A$ , called the *identity function on  $A$* , defined by

$$\text{id}_A(x) = x \quad \text{for all } x \in A.$$

**Example 2.33.** Most functions are neither injective nor surjective, for instance

The most fundamental operation on functions is *composition*: if  $f : A \longrightarrow B$  and  $g : B \longrightarrow C$ , then the composition  $g \circ f$  is the function  $A \longrightarrow C$  defined by

$$(g \circ f)(x) = g(f(x)) \quad \text{for all } x \in A.$$

If  $A = B$ , we have

If  $B = C$ , we have

**Definition 2.34.** Let  $A$  and  $B$  be sets. We say that a function  $f : A \longrightarrow B$  is *invertible* if there exists a function  $g : B \longrightarrow A$  such that

$$g \circ f = \text{id}_A \quad \text{and} \quad f \circ g = \text{id}_B.$$

**Theorem 2.35.** *Let  $f : A \longrightarrow B$  be a function from a set  $A$  to a set  $B$ . Then  $f$  is bijective if and only if it is invertible.*

## Orders

**Definition 2.36.** Let  $S$  be a set. A *(total) order* on  $S$  is a relation  $<$  satisfying the following two axioms:

(O1) for all  $x, y \in S$ , exactly one of the following statements is **True**:  $x < y$ ,  $y < x$ ,  $x = y$ ;

(O2) for all  $x, y, z \in S$ , if  $x < y$  and  $y < z$ , then  $x < z$ .

An *ordered set*  $S$  is a set with an order  $<$ .

We write  $x > y$  to mean

We write  $x \leq y$  and  $y \geq x$  to mean

**Example 2.37.**  $S = \mathbf{R}$  (or  $\mathbf{Q}$  or  $\mathbf{Z}$  or  $\mathbf{N}$ ) with the usual inequality relation  $<$  is an ordered set.

**Example 2.38.** A *singleton* (set with a single element)  $S = \{a\}$  is ordered via

**Example 2.39.** Let  $S = \{a, b, c\}$ . Which of the following relations is an order on  $S$ ?

- $a < b, a < c, b < c;$
- $b < a, c < a, b < c;$
- $a < b, b < c, c < a.$

**Theorem 2.40.** *Let  $S$  be an ordered set. If  $x, y \in S$  are such that  $x \leq y$  and  $y \leq x$ , then  $x = y$ .*

## 2.3 The real numbers

The most important sets in real analysis are arguably the subsets of  $\mathbf{R}$ , the set of real numbers.

This set has some interesting properties coming in two intermingling flavours:

- algebraically, it is a field under the usual operations of addition and multiplication;
- it has an order relation  $<$ .

Moreover,  $\mathbf{R}$  contains the field of rational numbers  $\mathbf{Q}$  in a way that is compatible with both the algebra and the order relation.

We will not dwell on the algebraic aspects: you are familiar with them, and they are better discussed in algebra subjects.

The properties of the order relation are crucial in analysis, so we will focus on these.

Some are given as axioms:

**(RO1)** for all  $x, y \in \mathbf{R}$ , exactly one of the following is **True**:

$$x < y, \quad y < x, \quad x = y;$$

**(RO2)** for all  $x, y, z \in \mathbf{R}$ , if  $x < y$  and  $y < z$  then  $x < z$ ;

**(RO3)** for all  $x, y, c \in \mathbf{R}$ , if  $x < y$  then  $x + c < y + c$ ;

**(RO4)** for all  $x, y \in \mathbf{R}$ , if  $0 < x$  and  $0 < y$  then  $0 < xy$ ;

**(RO5)** Completeness Axiom: every non-empty subset of  $\mathbf{R}$  that is bounded above in  $\mathbf{R}$  has a supremum in  $\mathbf{R}$ .

Other properties of the order relation can be proved from the axioms.

**Theorem 2.41.** *Let  $x, y, z \in \mathbf{R}$  with  $0 < z$ . If  $x < y$  then  $xz < yz$ .*

**Theorem 2.42.** *Let  $x, y \in \mathbf{R}$ . If  $x < y$  then  $-y < -x$ .*

**Theorem 2.43.** *Let  $x, y \in \mathbf{R}$ . If  $x < y$ , then*

$$x < \frac{x + y}{2} < y.$$

So far we have dealt with the order relation given by the strict inequality  $x < y$ . You should think about which of/whether the properties described above also hold for the non-strict inequality  $x \leq y$ .

We have not yet discussed the Completeness Axiom of  $\mathbf{R}$ . This requires a bit of terminology.

## 2.4 Bounds

Let  $S$  be an ordered set (e.g.  $S = \mathbf{R}$ ) and let  $A$  be a subset of  $S$ .

**Definition 2.44.** We say that  $A$  is *bounded below* in  $S$  if there exists  $\beta \in S$  such that

$$(\forall x \in A)\beta \leq x.$$

We say that  $\beta$  is a *lower bound* for  $A$  in  $S$ .

We say that  $A$  is *bounded above* in  $S$  if

We say that  $\beta$  is an *upper bound* for  $A$  in  $S$ .

We say that  $A$  is *bounded* in  $S$  if

**Example 2.45.** Let  $S = \mathbf{R}$  and consider the subset

$$A = \left\{ \frac{1}{r} : r \in \mathbf{R} \text{ and } r > 0 \right\}.$$

In the example, both  $-100$  and  $0$  are lower bounds for  $A$  in  $\mathbf{R}$ , but there is a clear sense in which  $0$  is more precise than  $-100$ .

Let  $S$  be an ordered set and let  $A$  be a subset of  $S$ .

**Definition 2.46.** Let  $L_A$  be the set of all lower bounds for  $A$  in  $S$ . We say that  $\alpha \in L_A$  is an *infimum* (or *greatest lower bound*) for  $A$  in  $S$  if

$$(\forall \beta \in L_A) \beta \leq \alpha.$$

Let  $U_A$  be the set of all upper bounds for  $A$  in  $S$ . We say that  $\alpha \in U_A$  is a *supremum* (or *least upper bound*) for  $A$  in  $S$  if

$$(\forall \beta \in U_A) \alpha \leq \beta.$$

**Theorem 2.47.** *Let  $A$  be a subset of an ordered set  $S$ .*

- *If a supremum for  $A$  in  $S$  exists, it is unique.*
- *If an infimum for  $A$  in  $S$  exists, it is unique.*

Because of this uniqueness property, it is unambiguous to write **sup**  $A$  for **the** supremum of  $A$ , and **inf**  $A$  for **the** infimum of  $A$ .

**Example 2.48.** Revisiting

$$A = \left\{ \frac{1}{r} : r \in \mathbf{R} \text{ and } r > 0 \right\},$$

**Example 2.49.** Consider

$$A = \{x \in \mathbf{R} : 1 \leq x < 2\}.$$

We can turn the argument in the previous example into a more general result:

**Theorem 2.50.** *Let  $A \subseteq \mathbf{R}$  be non-empty.*

- (a) Suppose  $A$  is bounded above and let  $\gamma \in \mathbf{R}$  be an upper bound for  $A$  in  $\mathbf{R}$ . Then  $\gamma = \sup A$  if and only if: for every  $\varepsilon > 0$  there exists  $x \in A$  such that  $x > \gamma - \varepsilon$ .*
- (b) Suppose  $A$  is bounded below and let  $\gamma \in \mathbf{R}$  be a lower bound for  $A$  in  $\mathbf{R}$ . Then  $\gamma = \inf A$  if and only if: for every  $\varepsilon > 0$  there exists  $x \in A$  such that  $x < \gamma + \varepsilon$ .*

We now have everything we need to make sense of the

**Completeness Axiom.** Every non-empty subset  $A \subseteq \mathbf{R}$  that is bounded above in  $\mathbf{R}$  has a supremum in  $\mathbf{R}$ .

The analogous statement for infimum also holds, and follows from the one for the supremum

There is an equivalent but more generalisable characterisation of completeness in terms of convergence of sequences, which we will discuss in the next chapter.

## The Archimedean Principle(s)

**Theorem 2.51** (Archimedean Principle I). *The subset  $\mathbf{N} \subseteq \mathbf{R}$  is not bounded above in  $\mathbf{R}$ .*

But wait, there's more!

**Theorem 2.52** (Archimedean Principle II). *For every  $x \in \mathbf{R}$  with  $x > 0$  there exists  $n \in \mathbf{N}$  such that  $n - 1 \leq x < n$ .*

**Theorem 2.53** (Archimedean Principle III). *If  $y, z \in \mathbf{R}$  with  $y, z > 0$  there exists  $n \in \mathbf{N}$  such that  $y < nz$ .*

**Theorem 2.54.** *The three variants of the Archimedean Principle are pairwise equivalent.*

## 2.5 Distance on the real line

An important class of subsets of  $\mathbf{R}$  consists of the *intervals*:

**Definition 2.55.** Given  $a, b \in \mathbf{R}$  with  $a \leq b$ , we have

- the *closed interval*  $[a, b] = \{x \in \mathbf{R} : a \leq x \leq b\}$ ;
- the *open interval*  $(a, b) = \{x \in \mathbf{R} : a < x < b\}$ ;
- $(a, b] = \{x \in \mathbf{R} : a < x \leq b\}$ ;
- $[a, b) = \{x \in \mathbf{R} : a \leq x < b\}$ ;
- $(a, \infty) = \{x \in \mathbf{R} : a < x\}$ ;
- $[a, \infty) = \{x \in \mathbf{R} : a \leq x\}$ ;
- $(-\infty, b) = \{x \in \mathbf{R} : x < b\}$ ;
- $(-\infty, b] = \{x \in \mathbf{R} : x \leq b\}$ .

Recall the *absolute value* function  $\mathbf{R} \rightarrow \mathbf{R}$  defined by

$$|x| = \begin{cases} -x & \text{if } x < 0, \\ x & \text{if } x \geq 0. \end{cases}$$

**Definition 2.56.** Given  $x, y \in \mathbf{R}$ , the *distance* between the points  $x$  and  $y$  on the real line is  $|x - y|$ .

**Theorem 2.57.** *Let  $x, a \in \mathbf{R}$  with  $a \geq 0$ . We have*

$$|x| \leq a \quad \text{if and only if} \quad -a \leq x \leq a,$$

*and similarly for the strict inequality  $|x| < a$ .*

**Theorem 2.58.** *For every  $x \in \mathbf{R}$  we have*

$$-x \leq |x| \quad \text{and} \quad x \leq |x|.$$

**Theorem 2.59** (The Triangle Inequality). *For every  $x, y \in \mathbf{R}$  we have*

$$|x + y| \leq |x| + |y|.$$

**Theorem 2.60.** *For every  $x, y \in \mathbf{R}$ , if  $x < y$  there exists  $r \in \mathbf{Q}$  such that  $x < r < y$ .*

**Corollary 2.61.** *For every  $z \in \mathbf{R}$  and every  $\varepsilon > 0$ , there exists  $r \in \mathbf{Q}$  such that  $|z - r| < \varepsilon$ .*

(A *corollary* is a theorem that follows easily from another theorem.)

A subset of  $\mathbf{R}$  with the property given in the Corollary is said to be *dense* in  $\mathbf{R}$ . So we have shown that  $\mathbf{Q}$  is dense in  $\mathbf{R}$ .